

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/321317769>

Assessment of prescribing practices at the primary healthcare facilities in Botswana with an emphasis on antibiotics: Findings and implications

Article in *International Journal of Clinical Practice* · November 2017

DOI: 10.1111/ijcp.13042

CITATIONS

31

READS

292

8 authors, including:



Yohana Mashalla

University of Botswana

66 PUBLICATIONS 989 CITATIONS

[SEE PROFILE](#)



Vincent Sethare

University of Botswana

9 PUBLICATIONS 51 CITATIONS

[SEE PROFILE](#)



Amos Massele

University of Botswana

33 PUBLICATIONS 762 CITATIONS

[SEE PROFILE](#)



Enoch Sepako

University of Botswana

62 PUBLICATIONS 443 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:




Disinvestment [View project](#)



CSA PerMed <http://www.permed2020.eu/> [View project](#)

ORIGINAL PAPER

Assessment of prescribing practices at the primary healthcare facilities in Botswana with an emphasis on antibiotics: Findings and implications

Yohana Mashalla¹ | Vincent Setlhare² | Amos Massele¹ | Enoch Sepako¹ |
Celda Tiroyakgosi³ | Joyce Kgatlwane⁴ | Mpo Chuma⁵ | Brian Godman^{6,7,8} 

¹Department of Biomedical Sciences, Faculty of Medicine, University of Botswana, Gaborone, Botswana

²Department of Family Medicine and Public Health, Faculty of Medicine, University of Botswana, Gaborone, Botswana

³Ministry of Health, Gaborone, Botswana

⁴School of Allied Health Professions, University of Botswana, Gaborone, Botswana

⁵School of Public Health, Faculty of Health Sciences, University of Botswana, Gaborone, Botswana

⁶Department of Clinical Pharmacology, Karolinska Institute, Stockholm, Sweden

⁷Strathclyde Institute of Pharmacy and Biomedical Sciences, Strathclyde University, Glasgow, UK

⁸Health Economics Centre, University of Liverpool Management School, Liverpool, UK

Correspondence

Brian Godman, Division of Clinical Pharmacology, Karolinska Institute, Karolinska University Hospital, Stockholm, Sweden.
Emails: brian.godman@ki.se;
brian.godman@strath.ac.uk

Funding information

University of Botswana Office of Research and Development

Summary

Background and Aims: Inappropriate drug prescribing has increased especially in developing countries where systems for monitoring medicine use are not well developed. This increases the rate of antimicrobial resistance. The study aim was to assess the prescribing patterns among urban primary health facilities in Botswana to provide future guidance including developing future quality indicators.

Methods: Retrospective data from patients' records between January and December 2013 in 19 clinics were collected in a cross-sectional study. The WHO/International Network for Rational Use of Drugs indicators were used to assess prescribing patterns in the study clinics.

Results: Average number of drugs per prescription was 2.8; 78.6% of the prescribed antibiotics were by International Non-proprietary Name and 96.1% complied with the Botswana Essential Drugs List. Overall rate of antibiotic prescribing was high (42.7%) with 14.7%, 5.9% and 1.3% of prescriptions having two, three and four antibiotics, respectively. Systemic antibiotics (JO1C) accounted for 45.4% of prescribed antibiotics of which amoxicillin accounted for 28.4% and metronidazole 14.4% of all antibiotic prescriptions. There was low use of co-amoxiclav (0.3% of all antibiotic prescriptions). Third generation cephalosporins and macrolides accounted for 9.8% and 6.2% of antibiotic prescriptions respectively, with no prescribing of fluoroquinolones. The majority of indications (87%) for antibiotic prescriptions were according to ICD classification.

Conclusions: While most indications for antibiotic prescriptions were based on signs and symptoms according to ICD, antibiotic prescribing rates were high with some conditions not requiring antibiotics because they are viral infections. There is a need to further improve prescribing practices through induction and training of in-service prescribers. An effective management tool for monitoring antibiotic prescribing practices at Primary Health Care facilities should be designed and implemented, including developing robust quality indicators.

1 | INTRODUCTION

The World Health Organization (WHO) defines the rational use of medicines as “patients receive medications appropriate to their clinical needs, in doses that meet their own individual requirements for an adequate period of time, at the lowest cost to them and their community”.¹ To address the irrational use of medicines, including inappropriate use of antibiotics leading to increasing rates of antimicrobial resistance (AMR) with its associated impact on morbidity, mortality and costs,²⁻⁸ the WHO in collaboration with the International Network for Rational Use of Drugs (INRUD) has developed core indicators for prescribing practices, patient care and facility-specific factors.^{2,9,10} These include the number of antibiotics per prescription without looking at issues of necessity or whether prescribing adheres to current guidelines.¹¹ The current WHO reference value for the average number of medicines per encounter is <2,^{10,12,13} with a comprehensive review between 1990 and 2009 reporting the average number among 104 countries was 2.6 for the Africa region.¹⁴ More recent reports indicate a 45.8% increase from 2.4 between 1995 and 2005 to 3.5 between 2006 and 2015.¹⁰ The percentage of medicines prescribed by the generic (International Non-proprietary Name; INN) name should be 100% (acceptable >80%).^{2,9,13} However, studies have reported lower rates for the African region at between 60% and 68%.^{5,12}

Currently, there is a scarcity of data regarding the quality of antibiotic prescribing in developing countries. In these settings, it is common for antibiotics to be prescribed for typically self-limiting conditions such as acute upper respiratory tract infections (URTIs) that are predominantly viral in origin and for other infections which do not require antibiotics.^{4,15-19} This situation is worse in African countries. The comparative report of antibiotic prescribing rates in the Africa region between 1995-2005 and 2006-2015 shows during the later period high antibiotic prescribing rates (22%, 58% and 22%) in Ghana, Tanzania and Nigeria, respectively,²⁰⁻²² with most of the increase because of prescribing in URTIs, which are typically viral in origin. Overall, approximately 50% of current antibiotic prescribing in outpatient departments is seen as inappropriate.²³ In Malaysia, as in Botswana, antibiotics are commonly and inappropriately prescribed by primary healthcare (PHC) providers at PHC facilities.²⁴⁻²⁷ Most of these reports though are based on small-scale studies, which could be underestimation of the extent of antibiotic use in developing countries.

Following the Alma Ata Declaration in 1978,²⁸ Botswana developed a successful and comprehensive PHC system with treatment provided in accordance with the standards set out in the Ministry of Health guidelines.²⁹ However to date, few studies have been undertaken to assess prescribing practices in PHC facilities in Botswana.³⁰⁻³² We have previously reported that 52 different documents/guidelines existed at 18 PHC facilities in the Greater Gaborone area of Botswana, of which 50% focused on treatment and management of diseases, the remaining 50% were general information and policy related.³³ Except for the guidelines for treating STIs found at 86% of the facilities, the majority of PHC facilities (56%) did not have current guidelines available and the Botswana Treatment Guide was only found at 50% of the clinics.³³

What's known

- There is considerable overuse of common antibiotics across countries particularly in primary healthcare centres, leading to increasing resistance rate.
- There are ongoing developments to improve the use of antibiotics across countries including WHO/International Network for Rational Use of Drugs (INRUD) indicators as well as guidelines.
- However, there are concerns with the specificity of WHO/INRUD indicators to fully assess the quality of antibiotic prescribing in ambulatory care as well as adherence to ICD codes when prescribing.

What's new

- There were high rates of International Non-proprietary Name prescribing in Botswana as well as high rates of prescribing of medicines contained in the Botswana EDL.
- However, there were high rates of antibiotic prescribing as well as high rates of prescribing two or more antibiotics on the first encounter. This may be because of a high prevalence of gynaecological and sexually transmitted infections in presenting patients in Botswana.
- The majority of indications for antibiotic prescribing (87%) were in accordance with ICD-10 codes; however rates could be improved to enhance future quality of prescribing

1.1 | Aims

The aim of this study was to assess current drug prescribing practices in PHC facilities in Gaborone and surrounding areas with a specific emphasis on antibiotic prescribing. The study results would be used to guide future corrective interventions. The interventions may include suggestions to develop new quality indicators, especially for antibiotic prescribing, in ambulatory care given the rather crude nature of current WHO/INRUD indicators.^{9,10}

2 | METHODOLOGY

This was a non-experimental cross-sectional descriptive study initially using the WHO/INRUD indicators⁹ before looking more specifically at antibiotic prescribing. The study was carried out between November and December 2015, collecting retrospective data from patient records for the period between January and December 2013. The PHC facilities were from Gaborone District, Tlokweng (South East District) and Mogoditshane (Kweneng District). The facilities are representative of urban settings in Botswana where the majority of patients seek their treatment from.

2.1 | Sample size and sampling procedures

The WHO⁹ recommends that there should be a minimum of 600 prescribing episodes included in a cross-sectional indicators survey, and where there are fewer than 20 facilities in a geographical or administrative region to be studied, all facilities should be included in the sample. The selected districts have a total of 20 PHC facilities; all were included in the study.

Data were collected by trained research assistants under the supervision of the principal researchers (YM and ES). Using patient's records, we estimated to collect at least 30 prescriptions from each PHC. To achieve this, patient prescription records were listed according to the month in which they were prescribed and assigned random numbers. Records were then randomly selected until a sample size of 30 patient records was realised. We wanted to assess initial prescriptions only in this study. Consequently, all records that showed re-attendances were excluded from the study. The diagnoses on the patient's records were used to determine the indication for antibiotic prescription without contact with the prescriber.

2.2 | Quality of antibiotic prescribing

The quality of antibiotic prescribing practices was assessed as follows: (i) prescribing against WHO/INRUD criteria^{9,10,13}; (ii) antibiotic prescribing against indicators developed and proposed by European organisations and (iii) antibiotic prescribing against the Botswana Essential Drug List.³⁴

The following indicators have been proposed and used by European organisations to assess the quality of systemic antibiotic (J01) prescribing³⁵⁻³⁷:

- Utilisation of penicillin (J01C) as a % of total antibiotic use.
- Utilisation of combination penicillin (eg, co-amoxiclav) as a % of total amoxicillin use.
- Utilisation of third- and fourth-generation cephalosporins vs first- and second- generation cephalosporins.
- Utilisation of fluoroquinolones (J01MA) as a % of total antibiotic use.

Prescribing rates at PHC facilities in Botswana were assessed against published rates among low and middle income European countries, including former Soviet Union Republics, as no baseline data currently exist for these facilities in Botswana.^{37,38}

The utilisation metric used is a prescription rather than internationally recognised metrics such as defined daily doses (DDDs) or DDDs per 1000 inhabitants per day (DIDs).³⁷⁻³⁹ This is because we wanted to assess antibiotic prescribing for the first indication, and we were evaluating antibiotic prescribing among providers at PHC facilities rather than the population as a whole.

2.3 | Data processing and analysis

For the purpose of improving the quality of data and the rigour of research results, data management procedures and processes were

in accordance with the methodology proposed by Vittinghoff et al⁴⁰. This included data preparation, cleaning, editing and creation of variables and identification of missing data. Since we were not testing any outcome variables against exposure interventions, and we were not assessing associations between the prescribers and prescribed drugs, only descriptive and inferential analysis frameworks were used to analyse prescribing information. Percentages, averages and frequencies were also used to describe the data.

2.4 | Ethical considerations

The study received ethical clearance from the University of Botswana, Institutional Review Board (Ref. URB/IRB/1506) and a permit to carry out the study in the PHC facilities was obtained from the Ministry of Health and Wellness (Ref. PME.13/18/87). The District Health Management Team Coordinator granted access to the facilities vide letter Ref: GGDHMT/14/2/i dated 28 November 2014. The identity of the clinics was protected by keeping them anonymous and assigned coded numbers.

3 | RESULTS

3.1 | General

Data from 19 PHC facilities only were analysed. Data from one clinic were not collected because during data collection, the clinic staff were on vacation. Out of an estimated 570 prescriptions that were collected, 20 prescriptions were excluded from the final analysis because seven patients had come to the clinics for routine Antenatal Clinic, six were patients who had visited the clinics for counselling services and another seven visited the facilities for HIV testing. This report is therefore based on 550 prescriptions.

3.2 | Prescribing practices against documented indicators

The total number of prescriptions analysed were 550 and the total number of medicines prescribed for all conditions were 1551, giving an average of 2.8 drugs per prescription. Of the total medicines prescribed, 1219 (78.6%) were prescribed using INN and 235 (42.7%) encounters contained at least one antibiotic prescription. 1490 (96.1%) were prescribed from the Botswana Essential Drug (EDL) (Table 1).

3.3 | Prescriptions with antibiotics

Table 2 shows that 306 antibiotics were prescribed during 235 patient encounters, of which 17 were topical applications (13 Chloramphenicol ointments, 3 Tetracycline ointments and one gentamicin ointment prescription). Systemic antibiotics (J01) were the most commonly prescribed. The beta-lactam antibiotics (J01C) accounted for 45.4% of prescriptions, rising to 48.1% when only systemic antibiotics were considered. Amoxicillin was the most commonly prescribed antibiotic

TABLE 1 Prescribing practices against documented indicators

Prescription practices	N	%
Total number of prescriptions analysed	550	
Total number of drugs prescribed	1551	
Average drugs/prescription	2.8	
Total prescriptions with generic (INN) names	1219	78.6
Total antibiotic encounters	235	42.7
Medicines from EDL	1490	96.1

(28.4%) followed by metronidazole (14.4%). Co-amoxiclav accounted for only 0.3% of all antibiotic prescriptions, while cotrimoxazole (sulphamethoxazole + tromethopim) accounted for 9.2% of all antibiotic prescriptions. Third generation cephalosporin (ceftriaxone) and macrolides (erythromycin) accounted for 9.8% and 6.2% of antibiotic prescriptions, respectively. This increased to 10.4% and 6.6%, respectively when only J01 systemic antibiotics were considered. No fluoroquinolones (J01MA) were prescribed.

3.4 | Indications for antibiotic prescriptions

Table 3 shows 69 diagnoses for which antibiotic prescriptions were written. Most diagnoses were based on signs and symptoms whilst some were specific disease conditions including asthma, diabetes, bronchitis, tonsillitis, conjunctivitis and chicken pox.

The majority of the antibiotic prescriptions (60) were compliant with the International Classification of Diseases (2017 ICD) while nine were not. The nine prescriptions were non-specific, unconventional and lacked crucial information such as sputum information on the amount, colour, odour and the presence of blood; "enlarged stomach" did not specify whether the enlargement was because of gas, solid mass, shifting or fluctuating; "local sepsis" did not contain information on location and nature of sepsis and for bacterial exudate the nature of the exudate (clear fluid, puss, odour) was not provided. Painful armpit lacked information on the nature of pain (throbbing, piercing, dull) and with or without induration.

Cough, vaginal discharge and sexually transmitted infections were most commonly indicated for antibiotic prescriptions in that order. For symptoms and signs involving the gastrointestinal system, diarrhoea was the commonest indication for antibiotic.

3.5 | Indications for more than one antibiotic per prescription

Out of the 235 encounters with antibiotic prescriptions, 45 (19.1%), 18 (7.7%) and 4 (1.7%) had two, three and four antibiotics per prescription, respectively. Doxycycline was the most common antibiotic combined as a second line drug followed by metronidazole and ceftriaxone, respectively (Table 4). The indications for combination of antibiotics were mostly sexually transmitted conditions including vaginal and urethral discharge and pelvic inflammatory diseases.

Antibiotic	ATC classification	Frequency (N)	Per cent
<i>Beta-lactams (J01C)</i>			
Amoxicillin	J01CA04	87	28.4
Ampicillin	J01CA01	3	1.0
Benzathine penicillin (Retarpen)	J01CE08	13	4.2
Cloxacillin	J01CF02	21	6.9
Co-amoxiclav	J01CR02	1	0.3
Crystalline penicillin	J01CE01	1	0.3
Penicillin V	J01CE02	13	4.2
<i>Total Beta-lactams</i>		139	45.4
Cotrimoxazole	J01EE01	28	9.2
Ceftriaxone	J01DD54	30	9.8
Chloramphenicol capsules	S01A01	3	1.0
Chloramphenicol ointment	S01AA01	13	4.2
Doxycycline	J01AA02	23	7.5
Erythromycin	J01FA01	19	6.2
Gentamicin	J01GB03	2	0.7
Metronidazole	J01XD01	44	14.4
Nitrofurantoin (nitrofurantoin derivative)	J01XE	1	0.3
Tetracycline ointment	S01AA	3	1.0
Gentamicin ointment	SO1AA	1	0.3
Total		306	100

TABLE 2 Commonly prescribed antibiotics at PHC facilities in Botswana

TABLE 3 Classification of the diseases, symptoms and signs indicated for antibiotics prescription at PHC facilities in Botswana by 2017 ICD10-CM/code

Condition	ICD-10-CM/code	Frequency
<i>General symptoms and signs</i>		
Fever	R50	10
Headache	R51	8
Dehydration	E86.0	1
<i>Symptoms and signs involving the circulatory and respiratory systems</i>		
Cough	R05	37
Acute tonsillitis	J03	18
Nasal congestion (common cold)	J00	7
Pharyngitis	J02	4
Sputum (unspecified)	R09.3	3
Swollen palate (acute laryngitis and tracheitis)	J04	2
Asthma	J45.909	2
Pleural thickening	L85.9	2
Upper respiratory infections	J06	2
Aspiration pneumonia	J69.0	1
Bronchitis	J20.0	1
<i>Symptoms and signs involving the digestive system and abdomen</i>		
Diarrhoea	R19	15
Abdominal and pelvic pain	R10	11
Loss of appetite	R63	6
Nausea and vomiting	R11	4
Constipation	K59.0	1
Enteritis	K52.9	1
Heartburn (epigastric pain)	R12	1
<i>Symptoms and signs involving the skin and subcutaneous tissues</i>		
Wound/sores	T81.30	12
Rash	CM/21	9
Boils and abscesses	CM/22	8
Skin condition unspecified	R23	7
Herpes zoster	B02	2
Chickenpox	B01	2
Burn	T30.0	2
Allergic dermatitis (unspecified)	L23.89	2
Acne	CM/706	1
Fibrotic	M72	1
Viral warts (genital)	B07	1
Implantation (sub-dermal contraceptive)	V25.5	1
Skin fungal infection (unspecified)	L08.9	1

(Continues)

TABLE 3 (Continued)

Condition	ICD-10-CM/code	Frequency
<i>Endocrine, nutritional and metabolic disease</i>		
Diabetes mellitus	E08	1
<i>Symptoms and signs involving genitourinary system</i>		
Vaginal discharge (unspecified)	N89.8	34
Sexually transmitted infections (unspecified)	A64	25
Dysuria (unspecified)	R30.0	6
Penile discharge (unspecified)	R36.9	5
Pelvic inflammatory disease	N73.9	4
Vagina itching (unspecified)	N89.4	3
Menstrual disorder syndrome	N92.6	3
Ovarian cyst	N83.2	3
Pregnancy masses	CM/09A	3
Orchitis	N45.2	2
Postsafe male circumcision (SMC)	Z41.2	2
Painful urinary bladder (unspecified)	N32.9	1
<i>Diseases of the ear and mastoid process</i>		
Otitis media	H60	7
Painful jaw		2
<i>Disorders of soft tissues and chest</i>		
Chest pain (unspecified)	R07.9	10
Backache	M54.9	2
Joint pain	M79.609	1
<i>Diseases of the eye and adnexa</i>		
Conjunctivitis	H10.0	5
Dacryocystitis (unspecified)	H04.309	1
<i>Symptoms and signs involving breast</i>		
Swollen breast (engorged)	N64.59	1
<i>Dental and oral diseases</i>		
Dental caries (unspecified)	K02.9	3
Chronic periodontitis	K05.3	2
Toothache (Unspecified)	K08.8	1
<i>Traumatic disorders</i>		
Injury (unspecified)	CM/929.9	14
Fractured index finger (unspecified)	S62.600A	1
<i>Other unspecified symptoms and signs</i>		
Altered discomfort		1
Swollen foreskin		1

(Continues)

TABLE 3 (Continued)

Condition	ICD-10-CM/code	Frequency
Enlarged stomach		1
Swellings (unspecified)		4
Local sepsis		2
Bacterial exudate		1
Itchy eyes		8
Painful armpit		1
Muscle pain (unspecified)		1

4 | DISCUSSION

When treatments were assessed against crude WHO/INRUD indicators, there was evidence of good prescribing practices in Botswana. This is shown by the fact that 96% of medicines prescribed were in accordance with the Botswana EDL and 79% of prescribing was by INN name (Table 2). The high rate of prescribing in accordance with the Botswana EDL³⁴ compares favourably with WHO recommendations of 100%^{10,12,13,41} as well as rates of 55%-80% reported for the Africa, Europe, Americas and South East Asia regions.^{2,10,42} The high prescribing rate from the EDL could be attributed to easy accessibility of the Botswana Treatment Guidelines at PHC facilities,³³ the routine availability of medicines incorporated into the Botswana EDL in PHCs, and the limited influence of pharmaceutical companies unlike other countries.^{2,21,43-46}

The high rate of INN prescribing in this study is comparable to 78% in the Western Pacific region but higher than rates of 27.7% and 48.9% reported in the Eastern Mediterranean and Southeast Asian regions⁴¹ as well as 68% for Africa as a whole.¹⁰ Possible explanations for high INN prescribing in this study could be a result of frequent in-service training workshops for health workers in Botswana sponsored by the government, benchmarking with colleagues in the region, the establishment of a medical school whose trainees are exposed to patient care early in their training where prescribing is typically INN, and easy access to digital information. This mirrors successful campaigns in other countries such as the UK with its high rate of INN prescribing.^{47,48} In order to attain WHO recommendation,^{9,10} we believe INN prescribing training should be intensified through regular training of in-service practitioners.

In this study, 42.7% of the encounters contained an antibiotic prescription, higher than <30% recommended by WHO.^{10,12,13} This is marginally higher than 39% and 34% for the Americas and European regions, but lower than 53% for the Eastern Mediterranean region⁴¹ and Africa as a whole (47%).¹⁰ This rate is also higher than 27% in a previous study in Botswana,³¹ and higher than 30% for Acute Respiratory Infections in children under 5 years old previously reported in Botswana.³² The high use of antibiotics could be because of a high burden of gynaecological and sexually transmitted infections (Table 3). There is also high prevalence (pandemic) of HIV in Botswana.⁴⁹ However, these patients are normally treated in specialist clinics, although in some cases, opportunistic infections may be treated in PHCs. The extent of multiple antibiotic prescribing is a

concern, but again may reflect the high prevalence of gynaecological and sexually transmitted infections in our study with up to three antibiotics recommended in the Botswana guidelines for these patients (Table 4). This will be a future area of research.

Regarding the quality of antibiotic prescribing, it is generally accepted as good clinical practice that antibiotics should be prescribed for specific diagnosis. We found that most of the indications for antibiotic prescriptions were based on signs and symptoms, rather than specific diagnoses, and were compliant with ICD (Table 3). However, nine conditions did not match the ICD which could either result from failure of patients to express themselves during history taking, poor history taking or the inability of prescribers to apply ICD codes when arriving at differential and definitive diagnoses. Adoption of ICD 10 codes for primary care diagnoses would go a long way towards creating a common language for classifying undifferentiated conditions that are seen in PHC settings.

There are concerns that some diagnoses such as a common cold, diarrhoea, painful legs, constipation and unclassified muscular pains, do not warrant antibiotics (Table 3). In addition, four prescriptions each containing antibiotics had no established diagnosis. The situation in this study is however better than reported in Nigeria where over 50% of the patient's folders reviewed had no established diagnosis.⁵⁰ The factors that contribute to over prescribing of antibiotics include inadequate in-service training, socio-cultural factors and patient demand.^{16,18,27,51} Most health facilities in Botswana and elsewhere are characterised by inadequate skilled human resources. These lower level healthcare workers are overwhelmed by the large number of patients seeking care and some pressurising on prescribers to prescribe injections and antibiotics.⁵²⁻⁵⁴ The establishment of the Medical School in 2009 is a dedicated effort by the Government of Botswana to improve the number and quality of healthcare providers in the country. Assessment of factors influencing antibiotic prescribing practices was outside the scope of this study but will be explored further in future research.

The high use of beta-lactamase antibiotics compares favourably with former Soviet Union Republics and Turkey with rates of 37.5%-65.6% of total J01 antibiotics.³⁷ The low use of co-amoxiclav is also encouraging as there are concerns that high utilisation increases side effects and resistance as well as costs.⁵⁵⁻⁵⁷ With rising concerns on the development of *Clostridium difficile* resistance,^{58,59} the low utilisation of cephalosporins in this study (Table 2) is also encouraging and compares favourably with rates of 0.5%-12.2% among former Soviet Union Republics and Turkey.³⁷ Similarly, the lack of fluoroquinolone prescribing is encouraging attributed to the non-inclusion of these medicines in the Botswana EDL.³⁴

In our future research, we will aim to explore adherence to treatment guidelines alongside establishing quality indicators for antibiotic prescribing among providers in both urban and rural PHC facilities in Botswana, which will assist prescribers with improving their quality of care.⁶⁰ Adherence is a complex phenomenon and typically far from optimal across countries.^{19,61-63} For instance, 46% of antibiotic prescriptions in a recent study were not indicated by the guidelines.⁶⁴ However, higher adherence rates were recently seen in Namibia (62%), although below national target rates of 95%.¹¹ Guideline adherence is influenced by several factors including the level of training

TABLE 4 The pattern of antibiotic combinations and the indications at PHC facilities in Gaborone, South East and Kweneng districts, Botswana

Antibiotics				Diagnoses	
Amoxicillin	Metronidazole			Cough	Chest pain
Amoxicillin	Gentamicin ear drops			Ear sores	
Amoxicillin	Metronidazole			Agisoris	Headache
Amoxicillin	Metronidazole	Retarpen		Pharyngitis	Swollen jaw
Amoxicillin	Cotrimoxazole			Post SMC	
Amoxicillin	Cotrimoxazole			Cough	Eye watering
Amoxicillin	Metronidazole			Abscess	
Amoxicillin	Ceftriaxone			PV discharge	
Amoxicillin	Metronidazole			Dental caries	
Augmentin	Doxycycline			Pelvic pain	Pleural thickening
Benzathine Penicillin	Doxycycline	Metronidazole	Ceftriaxone	PV discharge	
Benzathine Penicillin	Ceftriaxone			Gonococcal infection	
Benzathine Penicillin	Penicillin V			Tonsillitis	
Benzathine Penicillin	Amoxicillin			Sputum	
Benzathine Penicillin	Amoxicillin			NIL	
Benzathine Penicillin	Amoxicillin			Bacterial exudate	
Ceftriaxone	Metronidazole	Doxycycline		PV discharge	
Ceftriaxone	Doxycycline	Metronidazole		PV discharge	
Ceftriaxone	Doxycycline	Metronidazole		PV discharge	
Ceftriaxone	Doxycycline	Metronidazole		Penile rash	
Ceftriaxone	Metronidazole			Abdominal pain	
Ceftriaxone	Doxycycline			Urethral discharge	
Ceftriaxone	Doxycycline	Metronidazole		Pelvic pain	PV discharge
Ceftriaxone	Doxycycline	Metronidazole		PV discharge	
Ceftriaxone	Doxycycline	Metronidazole		STI discharge	
Ceftriaxone	Erythromycin			PV discharge	Gynaecological pelvis
Ceftriaxone	Doxycycline	Metronidazole		PV discharge	VRT
Ceftriaxone	Cotrimoxazole	Metronidazole	Doxycycline	PV discharge	
Ceftriaxone	Metronidazole	Erythromycin		Pregnant masses	
Ceftriaxone	Erythromycin	Metronidazole	Cotrimoxazole	PV discharge	Dysuria
Ceftriaxone	Doxycycline			Urethral discharge	
Cloxacillin	Bactrim			Abscess	
Cotrimoxazole	Ceftriaxone	Doxycycline	Metronidazole	PID	
Doxycycline	Ceftriaxone			Orchitis	
Doxycycline	Metronidazole			STI contact	
Doxycycline	Metronidazole			STI contact	
Doxycycline	Ceftriaxone	Metronidazole		PV discharge	
Gentamicin	Chloramphenicol			Dacryocystitis	Chest pain
Metronidazole	Ceftriaxone	Erythromycin		PV discharge	
Metronidazole	Ceftriaxone	Doxycycline		PV discharge	

Continues

of PHC personnel, available resources at healthcare facilities, workload, staff motivation, staff experience and the availability of managerial tools for monitoring prescribing patterns. Physicians may also misinterpret patient expectations, influencing adherence rates.⁶⁵

Effective monitoring can enhance adherence rates.⁶⁶ Care though must be taken in communicating appropriate guidelines, especially if there are differences between different policies and guidelines as recently seen in Namibia.⁶⁷

TABLE 4 Continued

Antibiotics				Diagnoses	
Amoxicillin	Metronidazole			Cough	Chest pain
Metronidazole	Cotrimoxazole			Abscess	Scapular wound and cold
Metronidazole	Doxycycline	Ceftriaxone		Pelvic pain	Ovarian cyst
Metronidazole	Doxycycline			MDS	PV discharge
Metronidazole	Erythromycin			Genital warts	Cough
Penicillin V	Cotrimoxazole			Tonsillitis	
Frequency	45	18	4		

MDS, menstrual disorder syndrome; PID, pelvic inflammatory disease; SMC, safe male circumcision; cold, common cold.

Based on our findings, we recommended that in order to improve prescribing at PHC facilities in Botswana, it is essential that prescribers be provided with in-service training on the use of current national treatment guidelines, classification of diseases, INN prescribing, as well as proper history taking and recording of diagnoses. In addition, Botswana should consider adopting ICD10 for primary care to aid easier classification of undifferentiated conditions that present in PHC facilities. Pharmacists should also play a key role in promoting rational prescribing practices by conducting drug utilisation reviews and using the findings to stimulate dialogue among practitioners to enhance the future rational use of medicines at PHC facilities.

The Ministry of Health and Wellness, as the major stakeholder responsible for policy formulation and guidance, should seek to design a harmonised managerial tool that will contain pertinent quality indicators for monitoring antibiotic prescribing practices at all PHC facilities. This includes targets for adherence to National Treatment Guidelines as well as classes of antibiotics prescribed, building on European and other guidance. Any quality indicators developed must reflect current conditions in ambulatory care as seen by the high prevalence of gynaecological and sexually transmitted infections in Botswana, which is not universal especially among European populations. This will help address concerns with current WHO/INRUD criteria. Similarly, multi-sector antimicrobial initiatives and programmes should be developed and implemented across all locations to improve future antibiotic prescribing through monitoring antimicrobial usage as well as developing and implementing strategic interventions aimed at optimising antimicrobial use and reducing AMR. This may help to improve antibiotic prescribing in the future in Botswana and reduce AMR rates. Hopefully any quality indicators developed will also be of interest to other African countries and wider with similar populations.

5 | LIMITATIONS

The study was carried out in an urban area of Botswana and did not include rural areas. However, as mentioned, this is where the majority of patients are currently treated in Botswana, although we recognise there could be differences in the characteristics of the respective patient populations. In addition, the results are from a retrospective collection of data, and information on the characteristics

of practitioners was not collected. However, we believe our findings will already give guidance to key stakeholders on ways to improve antibiotic prescribing in Botswana. We also believe our ongoing research to develop pertinent quality indicators across all locations, including both urban and rural PHCs, will be of interest to other African countries and wider.

ACKNOWLEDGEMENTS

This study was funded by the University of Botswana Office of Research and Development and the authors wish to acknowledge the Ministry of Health, Botswana for granting permission to carry out the study and to the staff and patients in the PHC facilities. We would also like to thank Dr Amanj Baker Kurdi for his helpful comments regarding potential statistical analyses.

DISCLOSURE

The authors declare no competing interests with this study.

AUTHOR CONTRIBUTIONS

YM, VS, AM, ES and MC designed the study and were involved in the collection and analysis. YH, AM and BG produced the initial draft manuscript. All authors critiqued successive drafts of the manuscript before submission and re-submission.

ORCID

Brian Godman  <http://orcid.org/0000-0001-6539-6972>

REFERENCES

1. WHO. The rational use of drugs. Report of a conference of experts, Nairobi, 25–29 November 1985. <http://apps.who.int/medicinedocs/documents/s17054e/s17054e.pdf>.
2. Ofori-Asenso R, Brhlikova P, Pollock AM. Prescribing indicators at primary health care centers within the WHO African region: a systematic analysis (1995–2015). *BMC Public Health*. 2016;16:724.
3. Fitchett JR, Atun R. Antimicrobial resistance: opportunity for Europe to establish global leadership. *Lancet Infect Dis*. 2016;16:388–389.

4. Oxford J, Kozlov R. Antibiotic resistance – a call to arms for primary healthcare providers. *Int J Clin Pract.* 2013;180(Suppl.):1-3.
5. Collignon P, Athukorala PC, Senanayake S, Khan F. Antimicrobial resistance: the major contribution of poor governance and corruption to this growing problem. *PLoS ONE.* 2015;10:e0116746.
6. WHO. Irrational drug use causing rise of anti-microbial resistance, 2005. <http://www.who.int/management/anmicrobialresistance.pdf>
7. Bhatt SP. Rational use of medicine. *Int J Pharm Res.* 2010;2:69.
8. WHO. World medicine situation report. Geneva, 2011. http://www.who.int/medicines/areas/policy/world_medicines_situation/WMS_ch6_wPricing_v6.pdf.
9. WHO. How to investigate drug use in health facilities: selected drug use indicators. EDM research series no. 00, 1993. <http://apps.who.int/medicinedocs/en/d/Js2289e/>
10. Ofori-Asenso R. Irrational use of medicines—a summary of key concepts. *Pharmacy.* 2016;4:35.
11. Nakwatumbah S, Kibuule D, Godman B, et al. Compliance to guidelines for the prescribing of antibiotics in acute infections at Namibia's national referral hospital: a pilot study and the implications. *Expert Rev Anti Infect Ther.* 2017;15:713-721.
12. Harvard Medical School and Harvard Pilgrim Health. Using indicators to measure country pharmaceutical situation fact book on WHO level I and level II monitoring indicators; 2006. <http://www.who.int/medicines/publications/WHOTCM2006.2A.pdf>
13. Dumoulin J, Kaddar M, Velasquez G. *Guide to the drug financing mechanisms.* Geneva: World Health Organization; 1998. <http://apps.who.int/medicinedocs/en/d/Jh2928e/>
14. Holloway KA, Ivanovska V, Wagner AK, Vialle-Valentin C, Ross-Degnan D. How we improved use of medicines in developing and transitional countries and do we know how to? Two decades of evidence. *Trop Med Int Health.* 2013;18:656-664.
15. Llor C, Bjerrum L. Antimicrobial resistance: risk associated with antibiotic overuse and initiatives to reduce the problem. *Ther Adv Drug Safety.* 2014;5:229-241.
16. Desta Z, Abula T, Beyene L, Fantahun M, Yohannes AG, Ayalew S. Assessment of rational drug use and prescribing in primary health care facilities in North West Ethiopia. *East Afr Med J.* 1997;74:758-763.
17. Olayemi SO, Akinyede AA, Oreagba AI. Prescription pattern at primary health care centres in Lagos. *Niger Postgrad Med J.* 2006;13:220-224.
18. Risk R, Naismith H, Brunett A, Moore SE, Cham M, Unger S. Rational prescribing in paediatrics in a resource-limited setting. *Arch Dis Child.* 2013;98:503-509.
19. van der Velden AW, Bell J, Sessa A, Duerden M, Altiner A. Sore throat: effective communication delivers improved diagnosis, enhanced self-care and more rational use of antibiotics. *Int J Clin Pract.* 2013; 67 (Suppl. 180):10-16.
20. Mohlala G, Peltzer K, Phaswana-Mafuya N, Namlagan S, Namlagan S. Drug prescription habits in public and private health facilities in 2 provinces in South Africa. *East Mediterr Health J.* 2010;16:324-328.
21. Bosu WK, Ofori-Adeji D. An audit of prescribing practices in health care facilities of the Wassa West District of Ghana. *West Afr J Med.* 2000;19:298-303.
22. Masele A, Nsimba S, Fulgence J. A survey of prescribing practices of health care workers in Kibaha District in Tanzania. *Tanz J Med.* 2007;22:31-33.
23. Dyar OJ, Beovic B, Vlahovic-Palcevski V, Verheij T, Pulcini C. How can we improve antibiotic prescribing in primary care? *Expert Rev Anti Infect Ther.* 2016;14:403-413.
24. Teng CL, Nik-Sherina H, Ng CJ, Chia YC, Atiya AS. Antibiotic prescribing for childhood febrile illness by primary care doctors in Malaysia. *J Paediatr Child Health.* 2006;42:612-617.
25. Teng CL, Tong SF, Khoo EM, et al. Antibiotics for URTI and UTI: prescribing in Malaysian primary care settings. *Aust Fam Physician.* 2011;40:325-329.
26. Mohd Fozi K, Kamaliah MN. The effect of profiling report on antibiotic prescription for upper respiratory tract infection. *Malays Fam Physician.* 2013;8:26-31.
27. Md Rezal RS, Hassali MA, Alrasheedy AA, Saleem F, Md Yusof FA, Godman B. Physician's knowledge, perceptions and behaviour towards antibiotic prescribing: a systematic review of the literature. *Expert Rev anti-Infect Ther.* 2015;13:665-680.
28. World Health Organisation. Declaration of Alma-Ata. International Conference on Primary Health Care, 1978. www.who.int/publications/almaata_declaration_en.pdf?ua=1
29. Ministry of Health. Botswana treatment guidelines 1992. MoH and National Standing Committee on Drugs, Gaborone.
30. Moffat HJ. *Prescribing practices study.* Gaborone: Ministry of Health; 1988.
31. Boonstra E, Linbaek M, Khulumani P, Ngome P, Fugelli P. Adherence to treatment guidelines in primary health care facilities in Botswana. *Trop Med Int Health.* 2002;7:178-186.
32. Boonstra E, Linbaek M, Ngome P. Adherence to management guidelines in acute respiratory infections and diarrhoea in children under 5 years old in primary health care in Botswana. *Int J Qual Health Care.* 2005;17:221-227.
33. Mashalla YJ, Sepako E, Sethlare V, Chuma M, Bulang M, Masele Y. Availability of guidelines and policy documents for enhancing performance of practitioners at the Primary Health Care (PHC) facilities in Gaborone, Tlokweng and Mogoditshane, Republic of Botswana. *J Publ Health Epid* 2016;8:127-135.
34. Ministry of Health. *The Botswana essential drug list* (2nd ed). 2012. <http://www.moh.gov.bw/>
35. Coenen S, Ferech M, Haaijer-Ruskamp FM, et al. European Surveillance of Antimicrobial Consumption (ESAC): quality indicators for outpatient antibiotic use in Europe. *Qual Saf Health Care.* 2007;16:440-445.
36. Adriaenssens N, Coenen S, Versporten A, Muller A, Vankerckhoven V, Goossens H. European Surveillance of Antimicrobial Consumption (ESAC): quality appraisal of antibiotic use in Europe. *J Antimicrob Chemother.* 2011;66(Suppl. 6):vi71-7
37. WHO. Antimicrobial Medicines Consumption (AMC) Network, 2017. <http://www.euro.who.int/en/publications/abstracts/antimicrobial-medicines-consumption-amc-network.-amc-data-20112014-2017>
38. Versporten A, Bolokhovets G, Ghazaryan L, et al. Antibiotic use in eastern Europe: a cross-national database study in coordination with the WHO Regional Office for Europe. *Lancet Infect Dis.* 2014;14:381-387.
39. WHO Collaborating Centre for Drug Statistics Methodology, Norwegian Institute of Public Health. Guidelines for ATC classification and DDD assignment 2010, 2011, 2012, 2013, 2014 and 2015. www.whocc.no/atc_ddd_publications/guidelines
40. Vittinghoff E, Glidden DV, Shiboski SC, McCulloch CE. *Regression methods in biostatistics: Linear, logistic, survival and repeated measures models.* New York, NY: Springer Science Business, Inc; 2005.
41. WHO, Harvard Medical School and Harvard Pilgrim Health (2009). Medicines use in primary care in developing and transitional countries Fact Book summarising results from studies reported between 1990 and 2006. http://www.who.int/medicines/publications/who_emp_2009.3/en/
42. WHO. Community-based surveillance of antimicrobial use and resistance in resource-constrained settings. Report from five pilot projects. <http://apps.who.int/medicinedocs/documents/s16168e/s16168e.pdf>
43. Masele AY, Nsimba SE, Rimoy G. Prescribing habits in church-owned primary health care facilities in Dar es Salaam and other Tanzanian coast regions. *East Afr Med J.* 2001;78:510-514.
44. Salmasi S, Ming LC, Khan TM. Interaction and medical inducement between pharmaceutical representatives and physicians: a meta-synthesis. *J Pharmaceut Pol Pract.* 2016;9:37.
45. Spurling GK, Mansfield PR, Montgomery BD, et al. Information from pharmaceutical companies and the quality, quantity, and

- cost of physicians' prescribing: a systematic review. *PLoS Med.* 2010;7:e1000352.
46. Fleischman W, Agrawal S, King M, et al. Association between payments from manufacturers of pharmaceuticals to physicians and regional prescribing: cross sectional ecological study. *BMJ.* 2016;354:i4189.
 47. Godman B, Bishop I, Finlayson AE, Campbell S, Kwon HY, Bennie M. Reforms and initiatives in Scotland in recent years to encourage the prescribing of generic drugs, their influence and implications for other countries. *Expert Rev Pharmacoecon Outcomes Res.* 2013;13:469-482.
 48. Godman B, Baker A, Leporowski A, et al. Initiatives to increase the prescribing of low cost generics; the case of Scotland in the international context. *Med Res Arch.* 2017;5:1-34.
 49. UNAIDS Gap Report, 2016. <http://www.avert.org/professionals/hiv-around-world/sub-saharan-africa/botswana>
 50. Isah H. Prescription patterns among care providers in Catholic-church-owned primary health care facilities in Northern Ecclesiastical provinces of Abuja, Jos and Kaduna, Nigeria. Preliminary findings. *J Pharm Allied Sci.* 2008. <http://dx.doi.org/10.4314/jophas.v5i2.48470>
 51. Kibuule D, Kagoya HR, Godman B. Antibiotic use in acute respiratory infections in under-fives in Uganda: findings and implications. *Expert Rev Anti Infect Ther.* 2016;14:863-872.
 52. Trap B, Hansen EH, Hogerzeil HV. Prescription habits of dispensing and non-dispensing doctors in Zimbabwe. *Health Policy Plan.* 2002;17:288-295.
 53. Tamuno I. Traditional medicine for HIV infected patients in antiretroviral therapy in a tertiary hospital in Kano, North-west Nigeria. *Asian Pac J Trop Med.* 2011;4:152-155.
 54. Rezal RS, Hassali MA, Alrasheedy AA, et al. Prescribing patterns for upper respiratory tract infections: a prescription-review of primary care practice in Kedah, Malaysia, and the implications. *Expert Rev Anti Infect Ther.* 2015;13:1547-1556.
 55. NHS Scotland. National Therapeutic Indicators 2014/ 2015. <http://www.sehd.scot.nhs.uk/publications/DC20141201nti.pdf>
 56. Desrosiers M, Evans GA, Keith PK, et al. Canadian clinical practice guidelines for acute and chronic rhinosinusitis. *Allergy Asthma Clin Immunol.* 2011;7:1-38.
 57. Appropriate use of amoxicillin clavulanate BPJ. 2011; 38: 28-33. http://www.bpac.org.nz/BPJ/2011/September/docs/bpj_38_amoxicillin_pages_28-33.pdf
 58. Dancer SJ. The problem with cephalosporins. *J Antimicrob Chemother.* 2001;48:463-478.
 59. NICE. *Clostridium difficile* infection: risk with broad spectrum antibiotics. <https://www.nice.org.uk/guidance/esmpb1/resources/clostridium-difficile-infection-risk-with-broad-spectrum-antibiotics-1502609568697285>
 60. Mainz J. Defining and classifying clinical indicators for quality improvement. *Int J Qual Health Care.* 2003;15:523-530.
 61. van Dijk L, de Jong JD, Westert GP, de Bakker DH. Variation in formulary adherence in general practice over time (2003-2007). *Fam Pract.* 2011;28:624-631.
 62. Shapiro DJ, Hicks LA, Pavia AT, Hersh AL. Antibiotic prescribing for adults in ambulatory care in the USA, 2007-09. *J Antimicrob Chemother.* 2014;69:234-240.
 63. Ivanovska V, Hek K, Mantel Teeuwisse AK, Leufkens HG, Nielen MM, van Dijk L. Antibiotic prescribing for children in primary care and adherence to treatment guidelines. *J Antimicrob Chemother.* 2016;71:1707-1714.
 64. Dekker AR, Verheij TJ, van der Velden AW. Inappropriate antibiotic prescription for respiratory tract indications: most prominent in adult patients. *Fam Pract.* 2015;32:401-407.
 65. Butler CC, Kinnersley P, Prout H, Rollnick S, Edwards A, Elwyn G. Antibiotics and shared decision-making in primary care. *J Antimicrob Chemother.* 2001;48:435-440.
 66. Bjorkhem-Bergman L, Andersen-Karlsson E, Laing R, et al. Interface management of pharmacotherapy. Joint hospital and primary care drug recommendations. *Eur J Clin Pharm.* 2013;69(suppl. 1):73-78.
 67. Kibuule D, Mubita M, Naikaku E, Kalemeera F, Godman BB, Sagwa E. An analysis of policies for cotrimoxazole, amoxicillin and azithromycin use in Namibia's public sector: Findings and therapeutic implications. *Int J Clin Pract.* 2017;71:e12918.

How to cite this article: Mashalla Y, Setlhare V, Massele A, et al. Assessment of prescribing practices at the primary healthcare facilities in Botswana with an emphasis on antibiotics: Findings and implications. *Int J Clin Pract.* 2017;71:e13042. <https://doi.org/10.1111/ijcp.13042>